

УДК 512.542

О P -СВОЙСТВЕ ПОДГРУПП КОНЕЧНЫХ ГРУПП

Баожун Ли, Амир Лиу

*Колледж прикладной математики, Ченгдуйский университет информационной технологии, Ченгду, Китай*ON P -PROPERTY OF SUBGROUPS OF FINITE GROUPS

Baojun Li, Aming Liu

College of Applied Mathematics, Chengdu University of Information Technology, Chengdu, China

Пусть H – подгруппа группы G . Мы говорим, что H имеет P -свойство в G , если $|G/K : N_{G/K}(HK/K \cap L/K)|$ является p -числом для любого pd -главного фактора L/K группы G . Используя это понятие, найдены некоторые новые критерии p -нильпотентности групп.

Ключевые слова: конечная группа, p -нильпотентная группа, P -свойство подгруппы.

Let H be a subgroup of a group G . We say that H has P -property in G if $|G/K : N_{G/K}(HK/K \cap L/K)|$ is a p -number for any pd -chief factor L/K of G . Using this property of subgroups, some new criterions of p -nilpotency of groups are obtained.

Keywords: finite group, p -nilpotent group, P -property of subgroup.

Introduction

Throughout this paper, all groups considered are finite and G always denotes a finite group. We use $\pi(G)$ to denote the set of all prime divisors of the order $|G|$ of G , π denotes a set of some primes and π' is the complement of π in the set \mathbb{P} of all primes. An integer n is called a π -number if all its prime divisors belong to π . G is said to be a π -group if $\pi(G) \subseteq \pi$. G is called a πd -group if $\pi(G) \cap \pi \neq \emptyset$. A class \mathfrak{F} of groups is called a formation if \mathfrak{F} is closed under taking homomorphic image and subdirect product. A formation \mathfrak{F} is said to be saturated if it contains every group G with $G/\Phi(G) \in \mathfrak{F}$. All unexplained notions and terminology are standard, as in [1], [2] or [3].

Recall that a subgroup A of G is said to permute with a subgroup B if $AB = BA$. It is known that AB is a subgroup of G if and only if A permutes with B . Thus the permutability of subgroups is very important. A subgroup H of G is called quasinormal [4] or permutable [3] in G if H permutes with all subgroups of G . If H permutes with all Sylow subgroups of G , then H is called s -permutable in G [5]. After the work in [5], [6], many authors attempted to study and apply other kinds of embedding properties of subgroups. For instance, a subgroup H in G is called seminormal [7], [8] if H is permutable with all subgroups of some supplement of H in G ; A subgroup H of G is called semipermutable in G [9] if it is permutable with every subgroup K of G with $(|H|, |K|) = 1$;

A subgroup H of a group G is said to be S -quasinormally embedded in G [10] if for each prime p dividing $|H|$, a Sylow p -subgroup of H is also a Sylow p -subgroup of some s -permutable subgroup of G ; A subgroup H of G is said to be conditionally permutable in G if H permutes with some conjugate of any subgroup of G [11]. More recently, W. Guo and A.N. Skiba [12] studied the structure of group by s -embedded and n -embedded subgroups.

It is known that if a p -group H is s -permutable in G then $|G : N_G(H)|$ is a p -number. Generalizing this property we give the following definition:

Definition 0.1. Let H be a subgroup of G . We call that H has P -property in G if for any pd -chief factor L/K of G , $|G/K : N_{G/K}(HK/K \cap L/K)|$ is a p -number.

In any finite soluble group G , it is easy to see that all permutable and s -permutable subgroups have P -property in G . Also, suppose that H is a p -subgroup of a group G (G is not necessary soluble), then H has P -property if H is seminormal, semipermutable, s -quasinormal embedded or so on in G (see proofs in Section 1).

The subgroups having P -property deeply influence the structure of a group. In this paper, we shall give some new criteria of p -nilpotency of groups.

2 Elementary Properties

In this section, we give some elementary results of P -property of subgroups, some of which are also lemmas of our main results.

Lemma 1.1. *Let H be a subgroup of G and N a normal subgroup of G . If H has P -property in G , then HN/N has P -property in G/N .*

Proof. (1) Let $(L/N)/(K/N)$ be a pd -chief factor of G/N . Then L/K is a pd -chief factor of G and hence $|G/K : N_{G/K}(HK/K \cap L/K)|$ is a p -number. It follows directly that

$$|(G/N)/(K/N):$$

$N_{(G/N)/(K/N)}((HK/N)/(K/N) \cap (L/N)/(K/N))|$ is a p -number and hence the lemma holds.

Let X be a subset of G , a subgroup H is said X -permutable with T if there is an element $x \in X$ such that $HT^x = T^xH$ (cf. [13]).

Proposition 1.2. *Let H be a p -subgroup of G . Then H has P -property in G if one of the following holds:*

- (1) H is normal in G ;
- (2) H is permutable in G ;
- (3) H is s -permutable in G ;
- (4) H is X -permutable with all Sylow subgroups of G , where X is a soluble normal subgroup of G .

Proof. Since all normal subgroups, all permutable subgroups and all s -permutable subgroups in G are X -permutable with all Sylow subgroups in G for any subgroup X of G , we only need to prove that H satisfies P -property in G when (4) holds. Let L/K be any pd -chief factor of G . Then HK/K is XK/K -permutable with all Sylow subgroups of G/K and XK/K is a soluble normal subgroup of G/K . If $K \neq 1$, then, by induction on $|G|$, we can assume that HK/K satisfies P -property in G/K and hence

$$|G/K : N_{G/K}(HK/K \cap L/K)|$$

is a p -number. Assume that $K=1$. Then L is minimal normal in G . Assume that L is abelian. Then L is a p -subgroup. If $H \cap L = 1$, then it holds clearly that $|G : N_G(H \cap L)| = 1$ is a p -number. Assume that $H \cap L \neq 1$. We claim that $|G : N_G(H \cap L)|$ is a p -number. Let q be any prime divisor of $|G|$ with $q \neq p$ and G_q a Sylow q -subgroup of G . Then there is an element $x \in X$ such that $HG_q^x = G_q^xH$. Since L is a normal p -group, we have that $L \cap HG_q^x = L \cap H$ is a normal subgroup of HG_q^x . Hence $G_q^x \subseteq N_G(H \cap L)$ and so $|G : N_G(H \cap L)|$ is a q' -number. Now, by the choice of q , we have obtained that $|G : N_G(H \cap L)|$ is a p -number when L is abelian. Assume that L is nonabelian. Then $L \cap X = 1$ and hence $X \subseteq C_G(L)$. Let q be any prime divisor of $|L|$ and Q a Sylow q -subgroup of L . Put G_q be a Sylow q -subgroup of G such that $Q \leq G_q$. By hypotheses, there is an element $x \in X$

such that $HG_q^x = G_q^xH$. Clearly,

$$(|HG_q^x : H|, |HG_q^x : G_q^x|) = 1$$

and $L \cap HG_q^x \leq HG_q^x$. By [2, Lemma 3.8.2],

$$\begin{aligned} L \cap HG_q^x &= (L \cap HG_q^x \cap H)(L \cap HG_q^x \cap G_q^x) = \\ &= (H \cap L)(L \cap G_q^x) = (H \cap L)Q^x. \end{aligned}$$

Since $X \subseteq C_G(L)$, $Q^x = Q$ and so $H \cap L$ permutes with Q . By the choice of Q , we see that $H \cap L$ is s -permutable in L and so is subnormal in L . Since L is minimal normal in G , L is a product of some simple groups. Thus $H \cap L \leq L$. Now consider that q is a prime different from p . Then $H \cap L$ char $(H \cap L)Q$ since $(H \cap L)Q \leq L$. On the other hand, $(H \cap L)Q = L \cap HG_q^x \leq HG_q^x$ by the above argument. Thus $G_q^x \subseteq N_G(H \cap L)$ and hence $|G : N_G(H \cap L)|$ is a q' -number for any $q \neq \pi(H \cap L)$. Therefore, $|G : N_G(H \cap L)|$ is a p -number and the proposition holds.

Proposition 1.3. *Let H be a p -subgroup of G . If H is seminormal or semipermutable in G , then H satisfies P -property in G .*

Proof. Assume that H is seminormal in G and let T be a supplement of H in G such that H permutes with all subgroup of T . Let q be any prime divisor of $|G|$ different from p and G_q an arbitrary Sylow q -subgroup of G . Then $G_q^x \subseteq T$ for some $x \in H$ since $G = HT$ and H is a p -subgroup. Thus H permutes with G_q^x and so $(HG_q^x)^x = HG_q^x$ is a subgroup of G . It follows that H permutes with G_q . Let L/K be any pd -chief factor of G . Clearly, HK/K is seminormal in G/K . If $K \neq 1$, then, by induction on $|G|$, we can assume that HK/K satisfies P -property in G/K and hence $|G/K : N_{G/K}(HK/K \cap L/K)|$ is a p -number. Assume that $K=1$. Then L is minimal normal in G . If L is abelian, then it can be obtained that $|G : N_G(H \cap L)|$ is a p -number by an argument as in Proposition 1.2. Assume that L is nonabelian. We claim that $H \cap L = 1$. Let Q be any Sylow q -subgroup of L and choose G_q to be a Sylow q -subgroup of G containing Q . Then $HG_q = G_qH$ and, by [2, Lemma 3.8.2],

$$L \cap HG_q = (L \cap H)(L \cap G_q) = (H \cap L)Q.$$

This induce that $H \cap L$ permutes with all Sylow q -subgroup of L . Since L is nonabelian, $(H \cap L)Q \neq L$ by Burnside $p^a q^b$ -Theorem. It follows from [14, Theorem 3] that there is a proper normal subgroup N of L such that $H \cap L \subseteq N$ or $Q \subseteq N$. But $Q \subseteq N$ is nonsense since Q is a Sylow

q -subgroup of L and L is a direct product of some nonabelian simple groups which are isomorphic to each other. Hence $H \cap L \subseteq N$. Repeat this argument, we can find finally that $H \cap L$ is subnormal in L . Thus $H \cap L \subseteq O_p(L) = 1$. It follows that $H \cap L \trianglelefteq G$ and, certainly $|G : N_G(H \cap L)|$ is a p -number. Similarly, one can prove that H satisfies P -property in G if H is semipermutable in G and the proposition holds.

Lemma 1.4. *Let H be a p -subgroup of G and N a minimal normal subgroup of G . Assume that H has P -property in G . If there is a Sylow p -sub-group G_p of G such that $H \cap N \trianglelefteq G_p$, then $H \cap N = N$ or 1 .*

Proof. Since H has P -property in G , $|G : N_G(H \cap N)|$ is a p -number. On the other hand, since $H \cap N \trianglelefteq G_p$, $|G : N_G(H \cap N)|$ is a p' -number. Thereby, $H \cap N$ is normal in G and it follows that $H \cap N = N$ or 1 since N is minimal normal in G and the lemma holds.

Lemma 1.5. *Let H be a p -subgroup of G for some prime divisor p of $|G|$ and assume that H has P -property in G . Then any G -chief factor L/K which does not avoid by H is a p -factor and hence is abelian.*

Proof. Assume that L/K is a G -chief factor which does not avoid by H . Then $(L/K)/(K/K)$ is a chief factor of G/K and does not avoid by HK/K . Since HK/K satisfies P -property in G/K by lemma 1.1, we can obtain that $L/K \cong (L/K)/(K/K)$ is a p -factor by induction on $|G|$ if $K \neq 1$. Assume that $K = 1$. Since H has P -property in G , $|G : N_G(H \cap L)|$ is a p -number. This induces that $G = G_p N_G(H \cap L)$, where G_p is a Sylow p -subgroup of G contained $H \cap L$. By [2, Lemma 3.4.9], $(H \cap L)^G \subseteq G_p$ and hence $(H \cap L)^G \subseteq O_p(G)$. It follows that $L \subseteq O_p(G)$ since $L \cap O_p(G) \supseteq H \cap L \neq 1$ and L is minimal normal in G . Thus the lemma holds.

Recall that a group is called a C_π -group if it has at least one Hall π -subgroups and all its Hall π -subgroups are conjugate.

Lemma 1.6. *Let G be a C_π -group and $p \notin \pi$. Assume that P is a Sylow p -subgroup of G . If every maximal subgroup of P (except one) has a π -closed supplement in G , then G is π -closed.*

Proof. It can be proved similar to Lemma 2.2 in [15] by choose Q to be a Hall π -subgroup of G .

2 Some Results

Theorem 2.1. *Let p be a minimal prime divisor of $|G|$, E be a normal subgroup of G such that G/E is p -nilpotent. Suppose that P is a Sylow*

p -subgroup of E . If there is a subgroup D of P with $1 < |D| < |P|$ such that every subgroup H in $\Sigma_D = \{H \subseteq P \mid |H| = |D|, \text{ or } |H| = 2|D| \text{ and } \exp(H) > 2 \text{ whenever } P \text{ is nonabelian } 2\text{-group and } |P : D| > 2\}$ either has a p -nilpotent supplement in G or P -property in G , then G is p -nilpotent.

Proof. Assume that the theorem does not hold and choose G to be a counter example of minimal order. We divide the proof into the following steps.

$$(1) O_{p'}(G) = 1.$$

If $O_{p'}(G) \neq 1$, then the hypotheses still hold on $G/O_{p'}(G)$ by Lemma 1.1. It follows from the choice of G that $G/O_{p'}(G)$ is p -nilpotent and so is G .

$$(2) O_p(E) = O_p(G) \cap E \neq 1.$$

Let N be a minimal normal subgroup of G with $N \subseteq E$. Then $p \mid |N|$ by (1). Let G_p be a Sylow p -subgroup of G with $P \subseteq G_p$. If there is a subgroup H of order $|D|$ such that $1 \neq H \cap N \trianglelefteq G_p$ and H has P -property in G , then N is a p -group by lemma 1.5 and hence $O_p(E) \neq 1$. Assume that any such subgroup H is not P -normal in G . Then H have a p -nilpotent supplement in G . Let P_1 be any maximal subgroup of G_p . If $P_1 \cap N \neq 1$, then there must be a subgroup $H \subseteq P \cap P_1$ of order $|D|$ with $1 \neq H \cap N \trianglelefteq G_p$. In fact, if $|D| < P_1 \cap N$, then we can choose H to be a subgroup of $P_1 \cap N$, otherwise, we can choose H with $P_1 \cap N \leq H \leq P \cap P_1$ since $|D| \leq |P \cap P_1|$. Thereby, H and so P_1 has a p -nilpotent supplement in G . If $N_p = G_p \cap N$, which is a Sylow p -subgroup of N , is not of order p , then, for every maximal subgroup P_1 of G_p , we have $P_1 \cap N = P_1 \cap N_p \neq 1$. By the above argument, every maximal subgroup P_1 of G_p has a p -nilpotent supplement in G . If $p > 2$, then G is soluble and so is a $C_{p'}$ -group. If $p = 2$, then G is a $C_{p'}$ -group by [16, Main Theorem]. Now, by Lemma 1.6, G is p' -closed and so is p -nilpotent, a contradiction. Assume that N_p is cyclic of order p . Then N is simple since N is minimal normal in G . But p is the minimal prime divisor of $|G|$ and so is of $|N|$. Thus N is soluble since N_p is cyclic. It follows that $|N| = p$ and hence $O_p(E) \supseteq N \neq 1$.

$$O_p(E) = O_p(G) \cap E \text{ is clear and (2) holds.}$$

(3) *Let N be a minimal normal subgroup of G with $N \subseteq O_p(E)$. Then $|N| = |D|$.*

If $|N| > |D|$, then there is a subgroup H of N

such that $|H|=|D|$ and $H \trianglelefteq G_p$. If H has P -property in G , then $H=H \cap N=N$ or 1 by lemma 1.4, a contradiction. Assume that H has a p -nilpotent supplement U in G . Then $G=HU=NU$ and it follows that $N \cap U \neq 1$. Clearly, $N \cap U \trianglelefteq G$, so $N \subseteq U$ and $G=U$ is p -nilpotent, a contradiction. Thus we can assume that $|N| \leq |D|$. If $|N| < |D|$, then, by lemma 1.1, we see that the hypotheses hold on G/N and, by the choice of G , G/N is p -nilpotent. If $N \subseteq \Phi(G)$, then G is p -nilpotent since the formation of all p -nilpotent group is saturated. Assume that $N \not\subseteq \Phi(G)$. Then N is complemented in G . Let X be a complement of N in G and Q a Hall p' -subgroup of $X \cap E$. Then, since

$$X \cap E \cong (X \cap E)N/N = E/N \leq G/N$$

is p -nilpotent, $NQ/N \text{ char } E/N \trianglelefteq G/N$ and so $NQ \trianglelefteq G$. By Frattini Argument,

$$G = (NQ)N_G(Q) = NN_G(Q).$$

Since $O_p(G)=1$, $N_G(Q) \neq G$. Hence $G_p \not\subseteq N_G(Q)$ and so $G_p \cap N_G(Q) < G_p$. Assume that P_1 is a maximal subgroup of G_p containing $G_p \cap N_G(Q)$. If $N \subseteq P_1$, then

$$G_p = G_p \cap NN_G(Q) = N(G_p \cap N_G(Q)) \subseteq NP_1 = P_1,$$

a contradiction. Thus $N \not\subseteq P_1$. Since P_1 is maximal in G_p , $N \cap P_1$ is maximal in N and $N \cap P_1 \trianglelefteq G_p$. Clearly, $P \cap P_1$ is maximal in P and

$$|N \cap P_1| < |N| < |D| \leq |P \cap P_1|,$$

so there is a subgroup H of P_1 such that $N \cap P_1 < H \leq P \cap P_1$ and $|H|=|D|$. Thus $N \cap P_1 = N \cap H$. If H has P -property in G , then, by lemma 1.4, $N \cap P_1 = N \cap H = 1$ or N . If $N \cap P_1 = N$, then $N \subseteq P_1$, a contradiction. Thus $N \cap P_1 = 1$. It follows that N is cyclic of order p and therefore, G is p -nilpotent since G/N is p -nilpotent and p is a minimal prime divisor of $|G|$. This contradicts to the choice of G . By hypotheses, H has a p -nilpotent supplement in G and so has a p -nilpotent supplement in E . Assume that $E=HU$ for some p -nilpotent subgroup U of E . Then U contains some conjugate of Q since E is clearly p -soluble and U contains some Hall p' -subgroup of E . Without loss of generality, we can assume that $Q \subseteq U$. Hence $U \subseteq N_G(Q)$. Since

$$E = HU = (P \cap P_1)U,$$

$$\begin{aligned} P &= P \cap E = P \cap (P \cap P_1)U = \\ &= (P \cap P_1)(P \cap U) = \\ &= (P \cap P_1)(P \cap N_G(Q) \cap U) \leq \\ &\leq (P \cap P_1)(P_1 \cap U) = P \cap P_1. \end{aligned}$$

It follows that $P \subseteq P_1$ and so $N \subseteq P \subseteq P_1$, a contradiction. This contradiction shows that (3) holds.

(4) N is cyclic and so $|D|=|N|=p$.

Assume that N is noncyclic. We claim that all minimal subgroup of E/N of order p either has p -nilpotent supplement in G or has P -property in G . Assume A/N is of order p and $A/N \subseteq E/N$. Clearly, A is noncyclic since N is. Thus there is a maximal subgroup H of A different from N . Therefore, $A/N = HN/N$ and $|H|=|N|=|D|$. If A/N has no p -nilpotent supplement in G/N , then clearly, H has no p -nilpotent supplement in G . Hence, by hypotheses, H has P -property in G . By lemma 1.1, $A/N = HN/N$ has P -property in G/N . Analogously, one can prove that if P is a nonabelian 2-group and $|P:D| > 2$, then every cyclic subgroup of P/N of order 4 either has a p -nilpotent supplement in G/N or has P -property in G/N . Thus, if N is noncyclic, then the hypotheses hold on G/N . Thereby, G/N is p -nilpotent by the choice of G . By a similar argument as in (3), one can prove that G is p -nilpotent. This contradicts to the choice of G and hence N is cyclic and $|N|=|D|=p$.

(5) $O_p(E) \leq Z_\infty(G)$.

Since p is the minimal divisor of G , it is equivalent to prove that every G -chief factor L/K in $O_p(E)$ is of prime order.

Assume that there exists a G -chief factor in $O_p(G)$ which is not of prime order. Then we can choose a G -chief factor L/K in $O_p(G)$ such that $|L/K|$ is not a prime but $|U/V|$ is a prime for all chief factor U/V of G with $U \subseteq P$ and $|U| < |L|$.

Let $W = \bigcap_{U \subseteq K} C_G(U/V)$, where U/V is a

G -chief factor. Then, by [3, A, (12.3)], all elements in W of p' -order act trivially on K since they act trivially on each G -chief factor of K . Let $C = C_G(K)$.

Assume $L \not\subseteq C$. If $L \subseteq KC$, then

$$L \cap C / K \cap C \cong L/K$$

is a chief factor of G . By the choice of L/K , $|L/K|=|L \cap C / K \cap C|$ is a prime, a contradiction.

If $L \not\subseteq KC$, then it is easily to see that

$$LC/K = L/K \times KC/K$$

and thereby, all p' -elements in C act trivially on L/K . It follows that all p' -elements in W act trivially on L/K . Hence $W \subseteq C_G(L/K)$. Since $G/W = G / \bigcap_{U \subseteq K} C_G(U/V)$ is an abelian group of exponent dividing $p-1$ and $W \subseteq C_G(L/K)$, $G/C_G(L/K)$ is an abelian group of exponent

dividing $p-1$. Since L/K is G -irreducible, L/K is of prime order by [17, I, Lemma 1.3], a contradiction.

Now assume that $L \subseteq C$. Then $K \subseteq Z(L)$. Let a, b be elements of order p in L . Suppose $p > 2$ or P is abelian. Then $(ab)^p = a^p b^p [b, a]^{\frac{p(p-1)}{2}} = 1$. Hence the product of elements of order p is still of order p and so $\Omega = \{a \in L \mid a^p = 1\}$ is a subgroup of L . If $\Omega \subseteq K$, then all elements of W with p' -order act trivially on every element of L of order p since they act trivially on K . It follows from [18, IV, Satz 5.12] that all elements in W of p' -order act trivially on L . Thus $W \subseteq C_G(L/K)$ and, as above argument, L/K is of prime order, a contradiction. If $\Omega \not\subseteq K$, then $L = \Omega K$. Choose an element a in $\Omega \setminus K$ such that $\langle a \rangle K / K \subseteq L/K \cap Z(G_p/K)$. Let $H = \langle a \rangle$. If H has a p -nilpotent supplement U in G , then HK/K has a p -nilpotent supplement UK/K in G/K . Thus

$$G/K = (HK/K)(UK/K) = (L/K)(UK/K).$$

Since L/K is minimal normal in G/K and is abelian, $L/K \cap UK/K = 1$ or $L/K \subseteq UK/K$ and $UK/K = G/K$. If $L/K \cap UK/K = 1$, then

$$\begin{aligned} |L/K| &= |G/K : UK/K| = \\ &= |HUK/K : UK/K| \leq |H| = p. \end{aligned}$$

It follows that L/K is cyclic of order p , which contradicts to the choice of L/K . If $L/K \subseteq UK/K = G/K$, then L/K is cyclic since L/K is minimal normal in G/K and $G/K = UK/K \cong U/U \cap K$ is p -nilpotent. Hence H has no p -nilpotent supplement in G . Since a is of order p , by the hypotheses and (4), H has P -property in G and so HK/K satisfies P -property in G/K . It follows from lemma 1.4 that

$$L/K = HK/K \cap L/K = HK/K$$

is cyclic, a contradiction. This contradiction shows that (5) holds.

(6) E is p -nilpotent.

Assume that E is not p -nilpotent. Then E is not a p -group and so $O_p(E) < E$. Let $R/O_p(E)$ be a G -chief factor with $R \leq E$. Then, clearly, R is not p -nilpotent. Let X be a minimal non- p -nilpotent subgroup of R . Then $X = A \rtimes B$, where A is a p -group of exponent p or 4 (when A is a nonabelian 2-group) and B is a p' -group. If $A \subseteq O_p(E)$, then B acts trivially on A by (5), a contradiction. Thus $A \not\subseteq O_p(E)$. Hence, if $p > 2$ or P is abelian, then there are elements of order p in $R \setminus O_p(E)$, and, if P is a nonabelian 2-group, then there are elements of order 2 or 4 in $R \setminus O_p(E)$. Assume that there is an element a of order p in $R \setminus O_p(E)$ and let

$H = \langle a \rangle$. Assume that H has a p -nilpotent supplement U in G . If $H \subseteq U$, then $G = U$ is p -nilpotent. If $H \not\subseteq U$, then $|G : U| = |H| = p$. Since p is the minimal prime divisor of $|E|$, $U \trianglelefteq G$. It follows that G is p -nilpotent since U is. Thus H has no p -nilpotent supplement in G . By hypotheses and (4), H has P -property in G . It follows that $HO_p(G)/O_p(G)$ has P -property in $G/O_p(G)$. Since $H \not\subseteq O_p(G)$, we have that

$$R/O_p(G) = HO_p(G)/O_p(G)$$

by Lemma 1.4. Thus $R = HO_p(G)$ is a p -group, a contradiction. Thus there is no element of order p in $R \setminus O_p(E)$. Hence $p = 2$ and there is an element a of order 4 in $R \setminus O_p(E)$ and $a^2 \in O_p(G)$. By a similar argument as above, we can get a contradiction and hence E must be p -nilpotent.

(7) *The final contradiction*

By (1) and (6), we have that E is a p -group and hence $E \leq Z_\infty(G)$ by (5). Again by (5), we see that G is p -nilpotent since G/E is. This is the final contradiction and the theorem holds.

By Theorem 2.1, we can obtain the following corollaries.

Corollary 2.2. *Let p be the minimal prime divisor of $|G|$ and E be a normal subgroup of G such that G/E is p -nilpotent. Let P be a Sylow p -subgroup of E . If every maximal subgroup of P either has a p -nilpotent supplement in G or has P -property in G , then G is p -nilpotent.*

Proof. If P is of order p , then E is clear p -nilpotent. By induction, we can assume that $O_p(G) = 1$. It follows that E is a p -group and so is of order p . By N/C -Theorem, $G/C_G(E)$ is isomorphic to a subgroup of $\text{Aut}(E)$ and hence is of order dividing $p-1$. Since p is the minimal prime divisor of $|G|$, we see that $G/C_G(E) = 1$ and so $E \leq Z(G)$. Thus G is p -nilpotent since G/E is. If P is not of order p , then its maximal subgroup is not trivial and hence G is p -nilpotent by Theorem 2.1.

Corollary 2.3. *Let p be the minimal prime divisor of $|G|$ and E be a normal subgroup of G such that G/E is p -nilpotent. Let P be a Sylow p -subgroup of E . If every subgroup of P of order 2 and 4 (if P is a nonabelian 2-group) either has a p -nilpotent supplement in G or P -property in G , then G is p -nilpotent.*

Proof. It can be proved similar as Corollary 2.2.

The following corollary is direct from Propositions 1.2, 1.3 and Theorem 2.1.

Corollary 2.4. *Let p be the minimal prime divisor of $|G|$ and E be a normal subgroup of G*

such that G/E is p -nilpotent. Let P be a Sylow p -subgroup of E . If there is a subgroup D of P with $1 < |D| < |P|$ such that for every subgroup H in $\Sigma_D = \{H \subseteq P \mid |H| = |D|, \text{ or } |H| = 2|D| \text{ and } \exp(H) > 2 \text{ whenever } P \text{ is nonabelian } 2\text{-group and } |P : D| > 2\}$, one of the following holds:

- (1) H is s -permutable in G ;
- (2) H is seminormal in G ;
- (3) H is semipermutable in G ;
- (4) H is X -permutable in G , where X is a soluble normal subgroup of G ;
- (5) H has a p -nilpotent supplement in G , then G is p -nilpotent.

REFERENCES

1. Shemetkov, L.A. Formations of Algebraic Systems / L.A. Shemetkov, A.N. Skiba. – Moscow : Nauka, 1989.
2. Guo, W. The theory of classes of groups / W. Guo. – Dordrecht : Kluwer Academic Publishers Group, 2000.
3. Doerk, K. Finite soluble groups / K. Doerk, T. Hawkes. – Berlin : Walter de Gruyter & Co., 1992.
4. Ore, O. Contributions to the theory of groups of finite order / O. Ore // Duke Math. J. – 1939. – Vol. 5, № 2. – P. 431–460.
5. Kegel, O. Sylow-Gruppen und Subnormalteiler endlicher Gruppen / O. Kegel // Math. Z. – 1962. – № 78. – P. 205–221.
6. Deskins, W. On quasinormal subgroups of finite groups / W. Deskins // Math. Z. – 1963. – № 82. – P. 125–132.
7. Su, X. Seminormal subgroups of finite groups / X. Su // J. Math. (Wuhan). – 1988. – Vol. 8, № 1. – P. 5–10.
8. Foguel, T. On seminormal subgroups / T. Foguel // J. Algebra. – 1994. – Vol. 165, № 3. – P. 633–635.

9. Wang, L. On s -semipermutable maximal and minimal subgroups of Sylow p -subgroups of finite groups / L. Wang, Y. Wang // Comm. Algebra. – 2006. – Vol. 34, № 1. – P. 143–149.

10. Ballester-Bolinches, A. Sufficient conditions for supersolubility of finite groups / A. Ballester-Bolinches, M.C. Pedraza-Aguilera // J. Pure Appl. Algebra. – 1998. – Vol. 127, № 2. – P. 113–118.

11. Guo, W. Conditionally permutable subgroups and supersolubility of finite groups / W. Guo, K.P. Shum, A.N. Skiba // Southeast Asian Bull. Math. – 2005. – № 29. – P. 493–510.

12. Guo, W. Finite groups with given s -embedded and n -embedded subgroups / W. Guo, A.N. Skiba // Journal of Algebra. – 2009. – Vol. 321, № 10. – P. 2843–2860.

13. Guo, W. X -permutable maximal subgroups of Sylow subgroups of finite groups / W. Guo, K.P. Shum, A.N. Skiba // Ukraïn. Mat. Zh. – 2006. – Vol. 58, № 10. – P. 1299–1309.

14. Kegel, O. Produkte nilpotenter gruppen / O. Kegel // Arch. Math. – 1961. – № 12. – P. 90C93.

15. Skiba, A. On weakly s -permutable subgroups of finite groups / A. Skiba // J. Algebra, Vol. 315, № 1. – P. 192–209.

16. Gross, F. Conjugacy of odd order Hall subgroups / F. Gross // Bull. London Math. Soc. – 1987. – Vol. 19, № 4. – P. 311–319.

17. Weinstein, M. Between nilpotent and solvable / M. Weinstein (ed.). – Washington : Polygonal Publ. House, N. J., 1982.

18. Huppert, B. Endliche Gruppen. I. / B. Huppert. – Berlin : Springer-Verlag, 1967.

The author were supported by the NNSF of P.R. China (Grant 11471055) and the Scientific Research Foundation of CUIT (Grant CSRF201008).

Поступила в редакцию 29.07.14.